

# Effectiveness of Fungicides for Management of Strawberry Diseases

Nicole Ward Gauthier, Extension Plant Pathologist

This guide is a decision-making tool to help growers select fungicides from different chemical classes (FRAC).<sup>1</sup>  
Additional information can be found in a number of UK Cooperative Extension Service publications, including ID-232, or by contacting county Extension agents.

FRAC Code Fungicide Group <sup>1</sup>	Fungicide	Chemical	Phomopsis Leaf Blight <i>Phomopsis obscurans</i>	Leaf Spot <i>Mycosphaerella fragariae</i>	Powdery Mildew <i>Sphaerotheca macularis</i>	Gray Mold <i>Botrytis cinerea</i>	Leather Rot <i>Phytophthora cactorum</i>	Anthracnose Fruit Rot or Crown Rot <i>Colletotrichum spp.</i>	Red Stele <i>Phytophthora fragariae</i>
1 Methyl Benzimidazole	Upwardly systemic. Fungicide resistance risk high. Tank mix with fungicides from a different fungicide group (FRAC) to prevent or delay resistance development. Do not mix with copper.								
	Topsin M~	thiophanate-methyl	++	++	+++	+++			
3 Sterol Inhibitors (DMI or SI)	Upwardly systemic. Rainfast in 2 hours. Some curative activity. There is wide variation in activity within this group. Medium risk for resistance. Make no more than 5 applications per season. Note: This group, which was formerly known as De-Mehtylation Inhibitors (DMI), are now classified as Sterol Biosynthesis Inhibitors (SBI or SI).								
	Orbit	propiconazole	+++	++	+++				
	Mettle	tetraconazole	+++	++	+++				
	Procure	triflumizole	+++		+++				
	Rally	myclobutanil	+++	++	+++				
	Quadris Top*	azoxystrobin + difenoconazole	+++	+++	+++	++	+++	+++	
4 Phenylamides (PA)	Systemic. Effective on water molds (oomycetes). There are various formulations and combinations of Ridomil fungicides on the market. Read labels carefully before use. Do not make more than 4 applications per season.								
	Ridomil	mefenoxam					+++		+++

FRAC Code Fungicide Group <sup>1</sup>	Fungicide	Chemical	Phomopsis Leaf Blight <i>Phomopsis obscurans</i>	Leaf Spot <i>Mycosphaerella fragariae</i>	Powdery Mildew <i>Sphaerotheca macularis</i>	Gray Mold <i>Botrytis cinerea</i>	Leather Rot <i>Phytophthora cactorum</i>	Anthracnose Fruit Rot or Crown Rot <i>Colletotrichum spp.</i>	Red Stele <i>Phytophthora fragariae</i>
7 Succinate Dehydrogenase Inhibitors (SDHI)	Upwardly systemic fungicide with varying degrees of systemic activity. Medium to high risk for resistance.								
	Pristine*	pyraclostrobin + boscalid	++	++	+++	++	+++	+++	
9 Anilino Pyrimidines (PA)	Locally systemic. Effective substitutes for Topsin M and Benlate for Botrytis fruit rot/gray mold. Fungicide timing is important. Medium risk for resistance development. Apply no more than 3 applications per season.								
	Fontelis	penthiopyrad			+++	+++			
	Scala	pyrimethanil				+++			
	Switch*	cyprodinil + fludioxonil	+			+++			
11 Quinone outside inhibitors (QoI)	Locally systemic. Fungicide resistance risk high. Do not make more than 4 applications per season. Note: These fungicides are also known as strobilurins.								
	Abound	azoxystrobin	++		+++	++	+++	+++	
	Cabrio	pyraclostrobin	++	++	+++	++	+++	+++	
	Pristine*	pyraclostrobin + boscalid	++	++	+++	++	+++	+++	
	Quadris Top*	azoxystrobin + difenoconazole	+++	+++	+++	++	+++	+++	
12 Phenyl pyrroles	Effective as good substitute for strobilurins in a resistance program. Low to medium risk for resistance. Do not make more than 4 applications per season.								
	Switch*	cyprodinil + fludioxonil				+++		++	
13 Azanaphthalenes	Upwardly systemic fungicide with some downward-moving ability. Effective protection against powdery mildew. Most effective during the early stages of powdery mildew development, as a protectant only. Medium risk for resistance. Do not make more than 5 applications per season.								
	Quintec	quinoxifen			+++				

FRAC Code Fungicide Group <sup>1</sup>	Fungicide	Chemical	Phomopsis Leaf Blight <i>Phomopsis obscurans</i>	Leaf Spot <i>Mycosphaerella fragariae</i>	Powdery Mildew <i>Sphaerotheca macularis</i>	Gray Mold <i>Botrytis cinerea</i>	Leather Rot <i>Phytophthora cactorum</i>	Anthracnose Fruit Rot or Crown Rot <i>Colletotrichum spp.</i>	Red Stele <i>Phytophthora fragariae</i>
17 SBI: Class III	Locally systemic. Low to medium risk for resistance. Maximum of 3 applications per season.								
	CaptEstate*	captan + fenhexamid	++	++		+++	+	++	++
	Elevate*	fenhexamid				+++			
33 Phosphonates	Fully systemic. Effective as a protectant treatment of water molds (oomycetes) such as <i>Phytophthora</i> spp.								
	Agri-Fos	phosphorus acid					+++		++
	Aliette	fosetyl-AL					+++		++
	Legion	fosetyl-AL					+++		++
	ProPhyt	phosphorus acid					+++		++
M Multi-site Inhibitors	Broad spectrum protectants. Best applied early-season as a protectant or as a contact fungicide. No systemic activity (washes off in the rain). Fungicide resistance risk low.								
	Captan	captan	++	++		++	+	++	+
	CaptEstate*	captan + fenhexamid	++	++		+++	+	++	++
	Thiram	thiram	++	++		++	+	++	+
U6 Phenyl-acetamide	Do not exceed 2 applications per season. Valued for late-season management of powdery mildew (3-day PHI).								
	Torino	cyflufenamid			+++				

\* Chemical contains more than one active ingredient, thus more than one FRAC code is assigned.

~ Never apply Topsin M alone. Combine with unrelated fungicide such as Captan or Thiram.

+++ highly effective

++ moderately effective

+ slightly effective

March 2016

**Check fungicide labels for specific host information, possible phytotoxicity, rates, re-entry intervals, and resistance management information. Always follow label instructions.**