

Black Knot

Nicole Ward Gauthier
Extension Plant Pathologist

Dennis Morgeson
County Extension Agent

INTRODUCTION

Black knot is a common, often serious, disease of plums and cherries in Kentucky. Ornamental *Prunus* species, as well as wild plums and cherries, may also be affected. Trees in both commercial and residential plantings are susceptible.

SYMPTOMS

Black knot is aptly named for the conspicuous black knotty growths that form on infected branches (FIGURE 1). Initially, however, these irregular swellings or knots are small and light brown (FIGURE 2). One year after infection, the enlarging knots become olive-green with a velvety surface. As the season progresses, swellings harden, become brittle, and turn black in color (FIGURE 3), reaching lengths of 6 inches by the end of the growing season. Often only one side of a limb is affected (FIGURE 4); but, in some cases, limbs may become completely encircled. Knots continue to expand each year until girdled branches eventually die.

FIGURE 1. BLACK KNOT DISEASE IS EASILY RECOGNIZED BY THE BLACK KNOTTY GROWTHS ON TWIGS AND BRANCHES.



CAUSE & DISEASE DEVELOPMENT

Black knot is caused by the fungus *Apiosporina morbosa* (syn. *Dibotryon morbosum*), which overwinters in knots on previously infected twigs and branches. Spores develop within knots in spring between bud break and shuck split and are spread by wind and rain. Only elongating (actively growing) twigs of the current season's growth are susceptible. While infection takes place in spring, knot development is not evident until autumn.



FIGURE 2



FIGURE 3




FIGURE 4

FIGURE 2. EARLY INFECTIONS APPEAR AS LIGHT BROWN SWELLINGS. **FIGURE 3.** KNOTS BECOME BLACK AND BRITTLE AS THE GROWING SEASON PROGRESSES. **FIGURE 4.** BLACK KNOT CANKERS OFTEN DEVELOP ON ONE SIDE OF A BRANCH, EXPANDING EACH YEAR.

DISEASE MANAGEMENT

Pruning and sanitation are the primary means for reducing or eliminating black knot. Fungicide applications may be used in conjunction with cultural practices, but are often not warranted.

Pruning

Pruning is best accomplished in autumn or winter after leaves fall. This makes it easier to locate branches with galls. Cuts should be made at the point of the branch union, at least 6 to 8 inches below knots. A second inspection in April (before buds open) is recommended to remove any newly formed knots. Complete removal of trees may be necessary if trunks and large limbs become girdled.

Sanitation

All diseased wood should be discarded and/or destroyed; avoid leaving infective clippings on the orchard floor. Removal of wild plum and cherry trees in the vicinity of cultivated trees can help reduce inoculum (infective spores).

Fungicides

If black knot is a persistent problem, fungicide sprays are recommended in spring to protect young, expanding twigs. Fungicides are not curative, and they are ineffective unless an eradication program of pruning and sanitation is followed. For specific recommended fungicides, homeowners should refer to Extension publication, *Disease and Insect Control for Home Grown Fruit in Kentucky* (ID-21), while commercial growers should refer to *Midwest Tree Fruit Spray Guide* (ID-92).

ADDITIONAL RESOURCES

- Commercial Tree Fruit Spray Guide (ID-92)
http://www2.ca.uky.edu/agcollege/plantpathology/ext_files/PPFShtml/MwTreeFruitSprayGuideID92.pdf
- Disease and Insect Control Program for Homegrown Fruit in Kentucky, including Organic Alternatives (ID-21)
<http://www.ca.uky.edu/agc/pubs/id/id21/id21.pdf>

Acknowledgement

The authors thank Paul Vincelli, UK Extension Plant Pathologist, for his review of this publication.

December 2015

Photos: University of Kentucky—Dennis Morgeson (figure 1), John Strang (figure 2) & Nicole Ward Gauthier (figure 4); Bugwood.org—Bruce Watt, University of Maine (figure 1 inset) & Ansel Oommen (figure 3)